

orion AGRISCIENCE



ECOTOXIC

KEEP OUT OF REACH OF CHILDREN

Carafe™

FUNGICIDE

For the control of powdery mildew and rust diseases on APPLES, STONEFRUIT, GRAPES, TAMARILLOS, PEAS, ORNAMENTALS and other VEGETABLES

Active Ingredient:

800 g/kg sulphur in the form of a water dispersible granule



4828



10 Kg



**HAZARD CLASSIFICATIONS: 6.4A, 9.1D****WARNING: HARMFUL, ECOTOXIC**

May cause eye irritation. Harmful to aquatic organisms.

PRECAUTIONS

STORAGE: Keep out of reach of children. Store in the original packaging in a secure place. Storage of certain quantities of this product trigger specific storage requirements (Refer to Safety Note). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of dust or spray mist. When handling, mixing or applying, wear overalls, chemical resistant boots and gloves, head covering and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

USE: Apply with well maintained and calibrated spray equipment.

DISPOSAL: Dispose of this product only by using in accordance with this label, or at a suitable landfill. Do not burn. Dispose of packaging to a suitable landfill. Do not burn. Do not use packaging for any other purpose.

ENVIRONMENTAL: Use this product carefully. Do not apply directly into or onto water. Do not allow spray drift outside the target area to occur. Take all reasonable steps to ensure that no significant adverse effects are caused to the environment beyond the application area.

First Aid

If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with water. If skin or hair contact occurs, remove contaminated clothing and flush with running water. For advice, call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately.

Emergency Response

In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 622)** (24 hrs).

Spillage

Exclude non-essential people from the area. Wear protective clothing. Moisten to prevent dusting and sweep up. Place in a waste container. Dispose of safely to a suitable landfill.

Safety Data Sheet

Additional information is provided in the safety data sheet. Freephone **0800 ORIONAS (0800 674 6627)** or visit **www.orionagriscience.co.nz**

GENERAL INFORMATION

Carafe is a protectant fungicide and must be applied before disease infection occurs. For best results, even spray coverage is necessary.

CAUTION: Some fruits and crops are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use on any crop unless local experience has proven sulphur to be safe in that locality. Do **NOT** apply Carafe when the temperature is above 25°C, or under conditions of extreme humidity, as crop damage can occur. Cucurbits and other crops grown under glass can be affected by an application of Carafe during periods of rapid growth.

Do **NOT** apply to sulphur shy apples such as Cox's Orange.

Do **NOT** apply to apricots.

Do **NOT** apply to pears.

Do **NOT** apply to apples with captan or within 5 days of a captan treatment.

Do **NOT** apply with crop oil or within 21 days of a crop oil application.

It is an offence to use this product on animals.

READ THE LABEL COMPLETELY BEFORE USE**DIRECTIONS FOR USE**

Crop	Disease	Rate per 100 Litres Water	Critical Comments
Apples	Powdery Mildew	100 - 200 grams	Apply from bud movement at 14 day intervals until harvest. Do NOT use post-blossom unless one pre-blossom sulphur application has been made. Carafe may affect fruit finish on some varieties.
Grapes	Powdery Mildew, Erineum Mite	150 - 300 grams	Apply from bud swell until 6 weeks before harvest at 10-14 day intervals. Use the higher rate where infection is established.
Ornamentals	Powdery Mildew, Rust.	100 - 200 grams	Apply at 10 - 14 day intervals when conditions are favourable to infection.
Peas	Powdery Mildew	100 - 200 grams	Apply when conditions are suitable for disease infection. Use the higher rate where infection has already occurred.
Nectarines, Peaches and Plums	Rust	100 - 200 grams	Apply at fruit set then at 21 day intervals until harvest. Use the higher rate during conditions of high pressure.
Tamarillos	Powdery Mildew	100 - 200 grams	Apply at 14 day intervals throughout the growing season. Use the higher rate if infection has occurred.
Vegetables. Do NOT apply to Cucumbers or rock melons	Powdery Mildew	100 - 200 grams	Use at regular 10 - 14 day intervals during periods of infection. Use the higher rate where infection has already occurred.

APPLICATION METHODS

Suitable for use with all applicators. The rates expressed are for high volume spraying to run off. For concentrated spraying, adjust dilution rate accordingly. Sulphur is corrosive so all spray equipment should be thoroughly flushed with clean water after use.

COMPATIBILITY

Carafe is compatible with most insecticides, fungicides and foliar fertilisers. When using with untried emulsifiable concentrate formulations, test the compatibility of both products by mixing small quantities before tank mixing. Do **NOT** use with spraying oils or any products containing mineral oil.

RAINFALL

Rain within 3 hours of treatment may reduce efficacy.

WITHHOLDING PERIODS

Nil.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P9169

See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSN0 Act 1996, No. HSR000741

See www.epa.govt.nz for approval conditions.

™CARAFE is a Trade Mark of Orion AgriScience Ltd.

Registrant: Agsin Pte Ltd, Zagro Global Hub, 5 Woodlands Terrace, Singapore 738430

Distributor: Orion AgriScience Ltd, Unit 1, 15 Sir Gil Simpson Drive, Harewood, Christchurch 8053
www.orionagriscience.co.nz

Batch No.

Date of Manufacture

orion
AGRISCIENCE

120