

**orion** AGRISCIENCE

**HARMFUL**  
KEEP OUT OF REACH OF CHILDREN

# PastureGibb<sup>®</sup>

## Liquid

# PLANT GROWTH REGULATOR

Stimulates additional PASTURE growth.

**Active Ingredient:**

32 g/litre of gibberellic acid in the form of a soluble concentrate.



9 421004 533300

**5 Litres**

## PLEASE READ THE LABEL COMPLETELY BEFORE USE

### PRECAUTIONS:

**STORAGE:** Keep out of reach of children. Store in the original, tightly closed container, in a secure place. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

**PERSONAL PROTECTION:** Avoid contact with eyes and skin. Avoid inhalation of spray mist.

**USE:** Apply with well maintained and calibrated spray equipment.

**ENVIRONMENTAL:** Use this product carefully. Do not allow spray drift outside the target area to occur.

**SAFETY DATA SHEET:** Additional information is provided in the safety data sheet. Freephone **0800 ORIONAS (0800 674 6627)** or visit [www.orionagriscience.co.nz](http://www.orionagriscience.co.nz)

### GENERAL INFORMATION

PastureGibb Liquid contains the naturally occurring gibberellic acid GA3. When applied under appropriate growing conditions, it will stimulate additional pasture growth. PastureGibb Liquid contains a component surfactant system – the addition of a separate wetting agent is not necessary.

### DIRECTIONS FOR USE

Use PastureGibb Liquid on a 3 – 4 weekly cycle. Application is best made in autumn and spring where the optimum soil temperature range of 6 - 13°C is likely to occur. In winter, PastureGibb Liquid can be applied in most districts apart from Canterbury, Otago, Southland, and in high and exposed areas in other parts of the country, where winter temperatures are likely to be too cold to allow for a growth response.

Apply 1 – 5 days after grazing, ensuring that there is some green leaf present to ensure foliar uptake. PastureGibb Liquid is best suited to rotationally grazed pastures. If set stocking, remove stock prior to application, and keep stock out for 3 – 4 weeks before grazing. Stock may be grazed sooner in the 3 – 4 week cycle if pasture is likely to become rank. Grazing earlier than 5 – 7 days post-treatment is likely to result in little or no benefit.

Soil moisture and fertiliser levels must be adequate to support increased pasture growth. Do not apply to rank pasture, ungrazed pasture, pasture less than 1 year old, or on paddocks containing livestock. As weeds also respond

to PastureGibb Liquid, avoid spraying paddocks with toxic weeds or high weed populations that may taint milk.

**Rate of Application:** 250 mL/ha. Apply in a minimum of 100 litres of water per hectare.

### RAINFALL

Do not spray if rainfall is expected within 2 hours of spraying.

### WITHHOLDING PERIOD

Not required.

### WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P8189  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

®PASTUREGIBB is a Registered Trade Mark of  
Orion AgriScience Limited

**Registrant:** Orion AgriScience Ltd,  
Unit 1, 15 Sir Gil Simpson Drive,  
Harewood, Christchurch 8053.

[www.orionagriscience.co.nz](http://www.orionagriscience.co.nz)

Batch No:

Date of manufacture:

PASTUREGIBB LIQUID

**orion**  
AGRISCIENCE