

orion AGRISCIENCE



ECOTOXIC

KEEP OUT OF REACH OF CHILDREN

TriumphTM
GEL
HERBICIDE

Ready to use weed killing gel for direct application onto cut stems.

Active Ingredient:

43 g/kg picloram as the potassium salt in the form of a gel.



9 421004 531993

10 Litres



PLEASE READ THE LABEL COMPLETELY BEFORE USE

HAZARD CLASSIFICATIONS: 6.5B, 6.9B, 9.2A

WARNING: ECOTOXIC

Harmful. May cause sensitisation from prolonged skin contact. May cause organ damage from repeated ingestion at high doses.

Selective herbicide – Very toxic to some plant species.

PRECAUTIONS

APPROVED HANDLER: Required if used by a commercial contractor (otherwise not required).

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure place.

PERSONAL PROTECTION: Avoid contact with eyes and skin. When applying, wear overalls, boots, and gloves. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

DISPOSAL: Dispose of this product only by using in accordance with this label, or at a suitable landfill. Dispose of container to a suitable landfill. Do not burn. Do not use container for any other purpose.

ENVIRONMENTAL: Use this product carefully. Do not use on weeds immediately next to desirable plants, or in areas where their roots may extend. Do not use where gel may be washed into the root zone of desirable plants.

First Aid

If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. Call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately for advice.

Emergency Response

In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

Spillage

Exclude non-essential people from the area. Wear protective clothing. Clean up by scraping and place in a waste container.

Safety Data Sheet

Additional information is provided in the safety data sheet. Freephone **0800 ORIONAS (0800 674 6627)** or visit www.orionagriscience.co.nz.

GENERAL INFORMATION

Effective killing of problem weeds (including shrubs and trees) is achieved by swabbing the cut stump with gel. Depending on the size of the unwanted plant, there is usually no resprouting after application, and the plant will die over the following weeks. Complete death of particularly large trees or shrubs may take several months. Repeat treatments will be necessary if regrowth occurs.

Although Triumph Gel can be applied at any time of year, better results will usually be achieved in spring - autumn. Avoid using when it is raining, or where rain is expected within 12 hours of treatment.

DIRECTIONS FOR USE

Trees, Woody Weeds and Shrubs, eg, barberry, broom, buddleia, cotoneaster, elder, gorse, hawthorn, matagouri, pine, poplar, privet, sweet briar, tutu, wattle, willow, woolly nightshade, etc. Cut the stem off near to the ground (2 – 10cm), and immediately apply gel 3 – 5 mm thick onto the cut stump. Alternatively, for larger shrubs and trees with a trunk diameter greater than 15 cm, 4 or more evenly spaced cuts, about 2 cm deep, can

be made with an axe or saw into the trunk near ground level. Apply the gel liberally into the cuts.

Scrambling Perennial weeds, eg, banana passionfruit, blue morning glory, greater bindweed, honeysuckle (Himalayan, Japanese), ivy (Cape, English, German, etc), jasmine, old mans beard, vines, etc. Cut the stem off near to the ground (2 – 10cm), and immediately apply gel 3 – 5 mm thick onto the cut stump. To minimise risks to any desirable tree the weed may be climbing, care should be taken to avoid the gel contacting the desirable tree. Damage may also occur to desirable trees should the gel be washed into the root zone, and alternative control methods should be considered.

Weeds with creeping or bulbous roots, eg, agapanthus, wild ginger, etc. Cut all shoots back to the root structure, and immediately apply gel 3 – 5 mm thick to the cut surface of the root structure. On plants such as wild ginger, where extensive root structures can occur between shoots, drill a hole approximately 10 mm diameter into every fourth structure from a shoot (to almost full depth), and fill with gel.

Not for use on food or animal feed crops.

WARRANTY

This formulation is warranted to contain the specified active ingredients within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Approved pursuant to the HSN0 Act 1996, No. HSR000552

See www.epa.govt.nz for approval conditions

™TRIUMPH is a Trade Mark of Orion AgriScience Ltd

Distributor: Orion AgriScience Ltd,
Unit 1, 15 Sir Gil Simpson Drive,
Harewood, Christchurch 8053
www.orionagriscience.co.nz

Batch No:

Date of manufacture:

TRIUMPH GEL 10L



AGRISCIENCE

